



Research Paper

## Elliptic Sombor energy of a graph

Saeid Alikhani<sup>1,\*</sup>, Nima Ghanbari<sup>1</sup>, Mohammad Ali Dehghanizadeh<sup>2</sup>

<sup>1</sup>Department of Mathematical Sciences, Yazd University, 89195-741, Yazd, I. R. Iran

<sup>2</sup>Department of Mathematics, National University of Skills (NUS), Tehran, I. R. Iran

**Academic Editor:** Ivan Gutman

**Abstract.** Let  $G$  be a simple graph with vertex set  $V(G) = \{v_1, v_2, \dots, v_n\}$ . The elliptic Sombor matrix of  $G$ , denoted by  $A_{ESO}(G)$ , is defined as the  $n \times n$  matrix whose  $(i, j)$ -entry is  $(d_i + d_j) \sqrt{d_i^2 + d_j^2}$  if  $v_i$  and  $v_j$  are adjacent and 0 for another cases. Let the eigenvalues of the elliptic Sombor matrix  $A_{ESO}(G)$  be  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$  which are the roots of the elliptic Sombor characteristic polynomial  $\prod_{i=1}^n (\rho - \rho_i)$ . The elliptic Sombor energy  $E_{ESO}$  of  $G$  is the sum of absolute values of the eigenvalues of  $A_{ESO}(G)$ . In this paper, we compute the elliptic Sombor characteristic polynomial and the elliptic Sombor energy for some graph classes. We compute the elliptic Sombor energy of cubic graphs of order 10 and as a consequence, we see that two  $k$ -regular graphs of the same order may have different elliptic Sombor energy.

**Keywords.** elliptic Sombor matrix, elliptic Sombor energy, elliptic Sombor characteristic polynomial, eigenvalues, regular graphs.

**Mathematics Subject Classification (2020):** 05C09, 05C90.

### 1 Introduction

Let  $G = (V, E)$  be a simple graph, with vertex set  $V(G) = \{v_1, v_2, \dots, v_n\}$ . If two vertices  $v_i$  and  $v_j$  of  $G$  are adjacent, then we use the notation  $v_i \sim v_j$ . For  $v_i \in V(G)$ , the degree of the vertex  $v_i$ , denoted by  $d_i$ , is the number of the vertices adjacent to  $v_i$ .

Let  $A(G)$  be adjacency matrix of  $G$  and  $\lambda_1, \lambda_2, \dots, \lambda_n$  its eigenvalues. These are said to be

\*Corresponding author (Email address: [alikhani@yazd.ac.ir](mailto:alikhani@yazd.ac.ir)).

Received 19 August 2024; Accepted 25 August 2024

First Publish Date: 25 August 2024

the eigenvalues of the graph  $G$  and to form its spectrum [9]. The energy  $E(G)$  of the graph  $G$  is defined as the sum of the absolute values of its eigenvalues

$$E(G) = \sum_{i=1}^n |\lambda_i|.$$

Details and more information on graph energy can be found in [21,22,24,29]. There are many kinds of graph energies, such as Randić energy [1,3,5,12,23], distance energy [33], incidence energy [4], matching energy [7,27] and Laplacian energy [11].

Sombor index is defined as  $SO(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2}$  (see [19]). More details on Sombor index can be found in [2,6,8,10,13,17,28,30–32,34]. Recently, in [20], Gutman introduced Sombor matrix of a graph  $G$  as  $A_{SO}(G) = (r_{ij})_{n \times n}$ , where

$$r_{ij} = \begin{cases} \sqrt{d_i^2 + d_j^2} & \text{if } v_i \sim v_j \\ 0 & \text{otherwise.} \end{cases}$$

The eigenvalues of  $A_{SO}(G)$  are denoted by  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ , and are said to form the Sombor spectrum of the graph  $G$ . The Sombor characteristic polynomial  $\phi_{SO}(G, \lambda)$  is

$$\phi_{SO}(G, \lambda) = \det(\lambda I - A_{SO}(G)) = \prod_{i=1}^n (\lambda - \rho_i),$$

and Sombor energy  $E_{SO}(G)$  is

$$E_{SO}(G) = \sum_{i=1}^n |\rho_i|.$$

We refer the reader to [16,18,25,26] for more details on Sombor energy.

In a recent paper (see [14]), the elliptic Sombor index of  $G$  is defined as

$$ESO(G) = \sum_{uv \in E(G)} (d_u + d_v) \sqrt{d_u^2 + d_v^2}.$$

Motivated by the definition of the Sombor matrix, we define the elliptic Sombor matrix as  $A_{ESO}(G) = (r_{ij})_{n \times n}$ , and

$$r_{ij} = \begin{cases} (d_i + d_j) \sqrt{d_i^2 + d_j^2} & \text{if } v_i \sim v_j \\ 0 & \text{otherwise.} \end{cases}$$

The eigenvalues of  $A_{ESO}(G)$  are denoted by  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ , and are said to form the elliptic Sombor spectrum of the graph  $G$ . The elliptic Sombor characteristic polynomial  $\phi_{ESO}(G, \lambda)$  is

$$\phi_{ESO}(G, \lambda) = \det(\lambda I - A_{ESO}(G)) = \prod_{i=1}^n (\lambda - \lambda_i),$$

and elliptic Sombor energy  $E_{ESO}(G)$  is

$$E_{ESO}(G) = \sum_{i=1}^n |\lambda_i|.$$

Two graphs  $G$  and  $H$  are said to be *elliptic Sombor energy equivalent*, or simply  $\mathcal{E}_{ESO}$ -equivalent, written  $G \sim H$ , if  $E_{ESO}(G) = E_{ESO}(H)$ . It is evident that the relation  $\sim$  of being  $\mathcal{E}_{ESO}$ -equivalence is an equivalence relation on the family  $\mathcal{G}$  of graphs, and thus  $\mathcal{G}$  is partitioned into equivalence classes, called the  $\mathcal{E}_{ESO}$ -equivalent. Given  $G \in \mathcal{G}$ , let

$$[G] = \{H \in \mathcal{G} : H \sim G\}.$$

We call  $[G]$  the equivalence class determined by  $G$ . A graph  $G$  is said to be *elliptic Sombor energy unique*, or simply  $\mathcal{E}_{ESO}$ -unique, if  $[G] = \{G\}$ .

A graph  $G$  is called  $k$ -regular if all vertices have the same degree  $k$ . One of the famous graphs is the Petersen graph which is a symmetric non-planar 3-regular graph of order 10.

There are exactly twenty one 3-regular graphs of order 10 [15]. In the study of elliptic Sombor energy, it is natural to investigate the elliptic Sombor characteristic polynomial and elliptic Sombor energy of cubic graphs of order 10 and check whether they recognized by their elliptic Sombor energy among other 3-regular graphs with the same order. We denote the Petersen graph by  $P$ .

In the next section we compute the elliptic Sombor energy of specific graphs. In Section 3, we study the elliptic Sombor energy of cubic graphs of order 10. As a consequence we show that the Petersen graph cannot be determined by its elliptic Sombor energy.

## 2 Elliptic Sombor energy of specific graphs

In this section, we study the elliptic Sombor characteristic polynomial and the elliptic Sombor energy for certain graphs. Here we compute the elliptic Sombor characteristic polynomial of paths and cycles.

**Theorem 2.1.** *For every  $n \geq 5$ , the elliptic Sombor characteristic polynomial of the path graph  $P_n$  satisfy:*

$$\phi_{ESO}(P_n, \lambda) = \lambda^2 \Lambda_{n-2} - 90\lambda \Lambda_{n-3} + 2025 \Lambda_{n-4},$$

where for every  $k \geq 3$ ,  $\Lambda_k = \lambda \Lambda_{k-1} - 128 \Lambda_{k-2}$  with  $\Lambda_1 = \lambda$  and  $\Lambda_2 = \lambda^2 - 8$ . Also the characteristic polynomial of  $P_2, P_3$  and  $P_4$  are  $\lambda^2 - 8, \lambda^3 - 90\lambda$  and  $\lambda^4 - 218\lambda^2 + 2025$  respectively.

*Proof.* It is easy to see that the characteristic polynomial of  $P_2$  is  $\lambda^2 - 8$ , Also for  $P_3$  is  $\lambda^3 - 90\lambda$  and for  $P_4$  is  $\lambda^4 - 218\lambda^2 + 2025$ . Now for every  $k \geq 3$  consider

$$M_k := \begin{pmatrix} \lambda & -4\sqrt{8} & 0 & \dots & 0 & 0 \\ -4\sqrt{8} & \lambda & -4\sqrt{8} & \dots & 0 & 0 \\ 0 & -4\sqrt{8} & \lambda & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \lambda & -4\sqrt{8} \\ 0 & 0 & 0 & \dots & -4\sqrt{8} & \lambda \end{pmatrix}_{k \times k},$$

and let  $\Lambda_k = \det(M_k)$ . One can easily check that  $\Lambda_k = \lambda\Lambda_{k-1} - 128\Lambda_{k-2}$ . Now consider the path graph  $P_n$ . Suppose that  $\phi_{ESO}(P_n, \lambda) = \det(\lambda I - A_{ESO}(P_n))$ . We have

$$\phi_{ESO}(P_n, \lambda) = \det \left( \begin{array}{c|cccccc|c} \lambda & -3\sqrt{5} & 0 & 0 & \dots & 0 & 0 & 0 \\ -3\sqrt{5} & & & & & & & 0 \\ 0 & & & & & & & 0 \\ 0 & & & & & & & 0 \\ \vdots & & & & M_{n-2} & & & \vdots \\ 0 & & & & & & & 0 \\ 0 & & & & & & & -3\sqrt{5} \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & -3\sqrt{5} & \lambda \end{array} \right)_{n \times n}.$$

So,

$$\begin{aligned} \phi_{ESO}(P_n, \lambda) &= \lambda \det \left( \begin{array}{c|c} M_{n-2} & \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \\ \hline 0 \dots 0 & -3\sqrt{5} \end{array} \middle| \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \right) \\ &\quad + 3\sqrt{5} \det \left( \begin{array}{c|cccc|c} -3\sqrt{5} & -4\sqrt{8} & \dots & 0 & 0 \\ 0 & & & & 0 \\ \vdots & & M_{n-3} & & \vdots \\ 0 & & & & -3\sqrt{5} \\ \hline 0 & 0 & \dots & -3\sqrt{5} & \lambda \end{array} \right). \end{aligned}$$

And so,

$$\begin{aligned} \phi_{ESO}(P_n, \lambda) &= \lambda \left( \lambda \Lambda_{n-2} + 3\sqrt{5} \det \left( \begin{array}{c|c} M_{n-3} & \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \\ \hline 0 \dots 0 & -4\sqrt{8} \end{array} \middle| \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \right) \right) \\ &\quad - 45 \det \left( \begin{array}{c|c} M_{n-3} & \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \\ \hline 0 \dots 0 & -3\sqrt{5} \end{array} \middle| \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \right) \end{aligned}$$

Hence,

$$\begin{aligned}\phi_{ESO}(P_n, \lambda) &= \lambda(\lambda\Lambda_{n-2} - 45\Lambda_{n-3}) \\ &\quad - 45 \left( \lambda\Lambda_{n-3} + 3\sqrt{5} \det \left( \begin{array}{c|c} M_{n-4} & \begin{matrix} 0 \\ \vdots \\ 0 \\ 0 \end{matrix} \\ \hline 0 \dots 0 & -4\sqrt{8} \end{array} \right) \right) \\ &= \lambda(\lambda\Lambda_{n-2} - 45\Lambda_{n-3}) - 45(\lambda\Lambda_{n-3} - 45\Lambda_{n-4}),\end{aligned}$$

and therefore we have the result.  $\square$

**Theorem 2.2.** For every  $n \geq 3$ , the elliptic Sombor characteristic polynomial of the cycle graph  $C_n$  satisfy:

$$\phi_{ESO}(C_n, \lambda) = \lambda\Lambda_{n-1} - 1024\Lambda_{n-2} + ((-1)^{n+1})(2)(8\sqrt{8})^n,$$

where for every  $k \geq 3$ ,  $\Lambda_k = \lambda\Lambda_{k-1} - 8\Lambda_{k-2}$  with  $\Lambda_1 = \lambda$  and  $\Lambda_2 = \lambda^2 - 512$ .

*Proof.* Similar to the proof of Theorem 2.1, for every  $k \geq 3$ , we consider

$$M_k := \begin{pmatrix} \lambda & -8\sqrt{8} & 0 & 0 & \dots & 0 & 0 & 0 & 0 \\ -8\sqrt{8} & \lambda & -8\sqrt{8} & 0 & \dots & 0 & 0 & 0 & 0 \\ 0 & -8\sqrt{8} & \lambda & -8\sqrt{8} & \dots & 0 & 0 & 0 & 0 \\ 0 & 0 & -8\sqrt{8} & \lambda & \dots & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \lambda & -8\sqrt{8} & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & -8\sqrt{8} & \lambda & -8\sqrt{8} & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & -8\sqrt{8} & \lambda & 0 \end{pmatrix}_{k \times k},$$

and let  $\Lambda_k = \det(M_k)$ . We have  $\Lambda_k = \lambda\Lambda_{k-1} - 512\Lambda_{k-2}$ . Suppose that  $\phi_{ESO}(C_n, \lambda) = \det(\lambda I - A_{ESO}(C_n))$ . We have

$$\phi_{ESO}(C_n, \lambda) = \det \left( \begin{array}{c|c} \lambda & -8\sqrt{8} \ 0 \ 0 \ \dots \ 0 \ 0 \ -8\sqrt{8} \\ \hline -8\sqrt{8} & \\ 0 & \\ 0 & \\ \vdots & \\ 0 & \\ 0 & \\ -8\sqrt{8} & \end{array} \begin{array}{c} \\ \\ \\ M_{n-1} \\ \\ \\ \end{array} \right)_{n \times n}.$$

So,

$$\begin{aligned} \phi_{ESO}(P_n, \lambda) &= \lambda \Lambda_{n-1} + 8\sqrt{8} \det \left( \begin{array}{c|ccc} -8\sqrt{8} & -8\sqrt{8} & 0 & \dots & 0 \\ \hline 0 & & & & \\ \vdots & & & & \\ 0 & & & & \\ \hline -8\sqrt{8} & & & & \end{array} \right) \\ &\quad + (-1)^{n+1} (-8\sqrt{8}) \det \left( \begin{array}{c|ccc} -8\sqrt{8} & & & & \\ \hline 0 & & & & \\ \vdots & & & & \\ 0 & & & & \\ \hline -8\sqrt{8} & 0 & \dots & 0 & -8\sqrt{8} \end{array} \right). \end{aligned}$$

Hence,

$$\begin{aligned} \phi_{ESO}(C_n, \lambda) &= \lambda \Lambda_{n-1} + 8\sqrt{8} \left( -8\sqrt{8} \Lambda_{n-2} + (-1)^n (-8\sqrt{8})^{n-1} \right) \\ &\quad + (-1)^{n+1} (-8\sqrt{8}) \left( (-8\sqrt{8})^{n-1} + (-1)^n (-8\sqrt{8}) \Lambda_{n-2} \right), \end{aligned}$$

and therefore we have the result.  $\square$

Now we consider to star graph  $S_n$  and find its elliptic Sombor characteristic polynomial and elliptic Sombor energy. We need the following Lemma.

**Lemma 2.3.** [9] If  $M$  is a nonsingular square matrix, then

$$\det \begin{pmatrix} M & N \\ P & Q \end{pmatrix} = \det(M) \det(Q - PM^{-1}N).$$

**Theorem 2.4.** For  $n \geq 2$ ,

(i) The elliptic Sombor characteristic polynomial of the star graph  $S_n = K_{1,n-1}$  is

$$\phi_{ESO}(S_n, \lambda) = \lambda^{n-2} \left( \lambda^2 - (n-1)(n^4 - 2n^3 + 2n^2) \right).$$

(ii) The elliptic Sombor energy of  $S_n$  is

$$E_{ESO}(S_n) = 2n \sqrt{(n-1)(n^2 - 2n + 2)}.$$

*Proof.* (i) One can easily check that the elliptic Sombor matrix of  $K_{1,n-1}$  is

$$A_{ESO}(S_n) = n \sqrt{n^2 - 2n + 2} \begin{pmatrix} 0_{1 \times 1} & J_{1 \times n-1} \\ J_{n-1 \times 1} & 0_{n-1 \times n-1} \end{pmatrix}.$$

We have  $\det(\lambda I - A_{ESO}(S_n)) =$

$$\det \begin{pmatrix} \lambda & -n\sqrt{n^2 - 2n + 2}J_{1 \times (n-1)} \\ -n\sqrt{n^2 - 2n + 2}J_{(n-1) \times 1} & \lambda I_{n-1} \end{pmatrix}.$$

Using Lemma 2.3,  $\det(\lambda I - A_{ESO}(S_n)) =$

$$\lambda \det(\lambda I_{n-1} - n\sqrt{n^2 - 2n + 2}J_{(n-1) \times 1} \frac{1}{\lambda}(n\sqrt{n^2 - 2n + 2}J_{1 \times (n-1)})).$$

Since  $J_{(n-1) \times 1}J_{1 \times (n-1)} = J_{n-1}$ , so

$$\begin{aligned} \det(\lambda I - A_{ESO}(S_n)) &= \lambda \det(\lambda I_{n-1} - \frac{n^2}{\lambda}(n^2 - 2n + 2)J_{n-1}) \\ &= \lambda^{2-n} \det(\lambda^2 I_{n-1} - n^2(n^2 - 2n + 2)J_{n-1}). \end{aligned}$$

On the other hand, the eigenvalues of  $J_{n-1}$  are  $n - 1$  (once) and  $0$  ( $n - 2$  times), the eigenvalues of  $n^2(n^2 - 2n + 2)J_{n-1}$  are  $(n - 1)(n^2)(n^2 - 2n + 2)$  (once) and  $0$  ( $n - 2$  times). Therefore

$$\phi_{ESO}(S_n, \lambda) = \lambda^{n-2} (\lambda^2 - (n - 1)(n^4 - 2n^3 + 2n^2)).$$

(ii) It follows from Part (i). □

We close this section by computing the elliptic Sombor characteristic polynomial of complete bipartite graphs and their Sombor energy.

**Theorem 2.5.** For natural number  $m, n \neq 1$ ,

(i) The elliptic Sombor characteristic polynomial of complete bipartite graph  $K_{m,n}$  is

$$\phi_{ESO}(K_{m,n}, \lambda) = \lambda^{m+n-2} (\lambda^2 - mn(m^2 + n^2)).$$

(ii) The elliptic Sombor energy of  $K_{m,n}$  is  $2(m + n)\sqrt{mn(m^2 + n^2)}$ .

*Proof.* (i) It is easy to see that the elliptic Sombor matrix of  $K_{m,n}$  is

$$(m + n)\sqrt{m^2 + n^2} \begin{pmatrix} 0_{m \times m} & J_{m \times n} \\ J_{n \times m} & 0_{n \times n} \end{pmatrix}.$$

Using Lemma 2.3 we have

$$\det(\lambda I - A_{ESO}(K_{m,n})) = \det \begin{pmatrix} \lambda I_m & -(m + n)\sqrt{m^2 + n^2}J_{m \times n} \\ -(m + n)\sqrt{m^2 + n^2}J_{n \times m} & \lambda I_n \end{pmatrix}.$$

So  $\det(\lambda I - A_{ESO}(K_{m,n})) =$

$$\det(\lambda I_m) \det(\lambda I_n - (m+n) \sqrt{m^2 + n^2} J_{n \times m} \frac{1}{\lambda} I_m (m+n) \sqrt{m^2 + n^2} J_{m \times n}).$$

We know that  $J_{n \times m} J_{m \times n} = m J_n$ . Therefore

$$\begin{aligned} \det(\lambda I - A_{SO}(K_{m,n})) &= \lambda^m \det(\lambda I_n - \frac{1}{\lambda} m(m^2 + n^2)(m+n)^2 J_n) \\ &= \lambda^{m-n} \det(\lambda^2 I_n - m(m^2 + n^2)(m+n)^2 J_n). \end{aligned}$$

The eigenvalues of  $J_n$  are  $n$  (once) and  $0$  ( $n-1$  times). So the eigenvalues of  $m(m^2 + n^2)(m+n)^2 J_n$  are  $mn(m^2 + n^2)(m+n)^2$  (once) and  $0$  ( $n-1$  times). Hence

$$\phi_{SO}(K_{m,n}, \lambda) = \lambda^{m+n-2} (\lambda^2 - mn(m^2 + n^2)(m+n)^2).$$

(ii) It follows from Part (i). □

### 3 Elliptic Sombor energy of $k$ -regular graphs

In this section we consider 2-regular and 3-regular graphs. As a beginning of this section, we have the following easy lemma:

**Lemma 3.1.** *Let  $G = G_1 \cup G_2 \cup G_3 \cup \dots \cup G_n$ . Then*

$$(i) \quad \phi_{ESO}(G) = \prod_{i=1}^n \phi_{ESO}(G_i).$$

$$(ii) \quad E_{ESO}(G) = \sum_{i=1}^n E_{ESO}(G_i).$$

As an immediate result of Lemma 3.1, we have the following results:

**Proposition 3.2.** *(i) If  $e = v_r v_{r+1} \in E(P_n)$ , then  $E_{ESO}(P_n - e) = E_{ESO}(P_r) + E_{ESO}(P_s)$ , where  $r + s = n$ .*

$$(ii) \quad \text{If } e \in E(C_n), (n \geq 3), \text{ then } E_{ESO}(C_n - e) = E_{ESO}(P_n).$$

(iii) *Let  $S_n$  be the star on  $n$  vertices and  $e \in E(S_n)$ . Then for any  $n \geq 3$ ,*

$$E_{ESO}(S_n - e) = E_{ESO}(S_{n-1}).$$

Now consider the 2-regulars. Every 2-regular graph is a disjoint union of cycles. By Theorem 2.2, we can find all the eigenvalues of elliptic Sombor matrix of cycle graphs. Therefore by Lemma 3.1, we can find elliptic Sombor characteristic polynomial and elliptic Sombor energy of 2-regular graphs. Before we continue, we need the following easy result that is the direct conclusion of the definition of elliptic Sombor energy:






Figure 1. Cubic graphs of order 10.

**Proposition 3.3.** If  $G$  is a  $k$ -regular graph of order  $n$ , then

$$E_{ESO}(G) = (2k)^n E_{SO}(G).$$

In [16], we showed that the Sombor energy of  $K_n$  is  $E_{SO}(K_n) = 2(n-1)^2\sqrt{2}$ . So by Proposition 3.3, we have the following result:

**Theorem 3.4.** For  $n \geq 2$ , The elliptic Sombor energy of  $K_n$  is

$$E_{ESO}(K_n) = 2^{n+1}(n-1)^{n+2}\sqrt{2}.$$

Now, we consider to the elliptic characteristic polynomial of 3-regular graphs of order 10. Also we compute the elliptic Sombor energy of this class of graphs. There are exactly 21 cubic graphs of order 10 given in Figure 1 (see [15]). We have the Sombor energy of these graphs in table 1 as we computed them in [16]:

$G_i$	$E_{SO}(G_i)$	$G_i$	$E_{SO}(G_i)$	$G_i$	$E_{SO}(G_i)$
$G_1$	64.161	$G_8$	64.161	$G_{15}$	62.767
$G_2$	63.043	$G_9$	64.981	$G_{16}$	59.396
$G_3$	62.880	$G_{10}$	61.399	$G_{17}$	67.882
$G_4$	57.336	$G_{11}$	62.375	$G_{18}$	57.517
$G_5$	60.638	$G_{12}$	67.882	$G_{19}$	66.096
$G_6$	63.403	$G_{13}$	61.000	$G_{20}$	59.396
$G_7$	63.969	$G_{14}$	65.835	$G_{21}$	50.911

Table 1. Sombor energy of cubic graphs of order 10.

Now, by Using Table 1 and Proposition 3.3, we have the elliptic Sombor energy of cubic graphs of order 10 up to three decimal places, as we see in Table 2:

$G_i$	$E_{ESO}(G_i)$	$G_i$	$E_{ESO}(G_i)$	$G_i$	$E_{ESO}(G_i)$
$G_1$	13858.776	$G_8$	13858.776	$G_{15}$	13557.672
$G_2$	13617.288	$G_9$	14035.896	$G_{16}$	12829.536
$G_3$	13582.080	$G_{10}$	13262.184	$G_{17}$	14662.512
$G_4$	12384.576	$G_{11}$	13473.000	$G_{18}$	12423.672
$G_5$	13097.808	$G_{12}$	14662.512	$G_{19}$	12980.736
$G_6$	13695.048	$G_{13}$	13176.000	$G_{20}$	12829.536
$G_7$	13758.336	$G_{14}$	14220.360	$G_{21}$	10996.776

Table 2. Elliptic Sombor energy of cubic graphs of order 10.

**Proposition 3.5.** Six cubic graphs of order 10 are not  $\mathcal{E}_{ESO}$ -unique.

*Proof.* By Table 2, we see that  $[G_1] = \{G_1, G_8\}$ ,  $[G_{12}] = \{G_{12}, G_{17}\}$  and  $[G_{16}] = \{G_{16}, G_{20}\}$ . Therefore, we have fifteen cubic graphs of order 10 which are  $\mathcal{E}_{SO}$ -unique.  $\square$

As an immediate result of Proposition 3.5, we have:

**Corollary 3.6.** In general, two  $k$ -regular graphs of the same order may have different elliptic Sombor energy.




Figure 2. Petersen graph

**Theorem 3.7.** Let  $\mathcal{G}$  be the family of 3-regular graphs of order 10. For the Petersen graph  $P$  (Figure 2 or  $G_{17}$  in Figure 1), we have the following properties:

- (i) The Petersen graph  $P$  is not  $\mathcal{E}_{ESO}$ -unique in  $\mathcal{G}$ .
- (ii) The Petersen graph  $P$  has the maximum elliptic Sombor energy in  $\mathcal{G}$ .

*Proof.* (i) The Sombor matrix of  $P$  is

$$A_{ESO}(P) = \begin{pmatrix} 0 & 18\sqrt{2} & 0 & 0 & 18\sqrt{2} & 18\sqrt{2} & 0 & 0 & 0 & 0 \\ 18\sqrt{2} & 0 & 18\sqrt{2} & 0 & 0 & 0 & 18\sqrt{2} & 0 & 0 & 0 \\ 0 & 18\sqrt{2} & 0 & 18\sqrt{2} & 0 & 0 & 0 & 18\sqrt{2} & 0 & 0 \\ 0 & 0 & 18\sqrt{2} & 0 & 18\sqrt{2} & 0 & 0 & 0 & 18\sqrt{2} & 0 \\ 18\sqrt{2} & 0 & 0 & 18\sqrt{2} & 0 & 0 & 0 & 0 & 0 & 18\sqrt{2} \\ 18\sqrt{2} & 0 & 0 & 0 & 0 & 0 & 0 & 18\sqrt{2} & 18\sqrt{2} & 0 \\ 0 & 18\sqrt{2} & 0 & 0 & 0 & 0 & 0 & 0 & 18\sqrt{2} & 18\sqrt{2} \\ 0 & 0 & 18\sqrt{2} & 0 & 0 & 18\sqrt{2} & 0 & 0 & 0 & 18\sqrt{2} \\ 0 & 0 & 0 & 18\sqrt{2} & 0 & 18\sqrt{2} & 18\sqrt{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 18\sqrt{2} & 0 & 18\sqrt{2} & 18\sqrt{2} & 0 & 0 \end{pmatrix}.$$

So

$$\phi_{SO}(P, \lambda) = \det(\lambda I - A_{SO}(P)) = (\lambda - 9\sqrt{2})(\lambda + 6\sqrt{2})^4(\lambda - 3\sqrt{2})^5.$$

Therefore we have:

$$\lambda_1 = 1994\sqrt{2}, \quad \lambda_2 = \lambda_3 = \lambda_4 = \lambda_5 = -1296\sqrt{2}, \quad \lambda_6 = \lambda_7 = \lambda_8 = \lambda_9 = \lambda_{10} = 648\sqrt{2},$$

and so we have  $E_{SO}(P) = 10368\sqrt{2}$ . By Table 2, we have  $P \in \{G_{12}, G_{17}\}$ . Therefore  $P$  is not  $\mathcal{E}_{SO}$ -unique in  $\mathcal{G}$ .

(ii) It follows from Part (i) and Table 2. □

In [16], we have shown that if two connected  $k$ -regular graphs have the same Sombor energy, then their adjacency matrices may have or have not the same permanent. Now by Proposition 3.3, we have the following result:

**Proposition 3.8.** *If two connected  $k$ -regular graphs have the same elliptic Sombor energy, then their adjacency matrices may have or have not the same permanent.*

Also in [16], we have shown that if two graphs have the same permanent, then we can not conclude that they have same Sombor energy. Consequently, if we have two graphs with the same permanent, then we can not conclude that they have same elliptic Sombor energy.

We think that the elliptic Sombor energy of no graph is integer. We end this section with the following conjecture:

**Conjecture 3.9.** *There is no graph with integer-valued elliptic Sombor energy.*

## 4 Conclusions

In this paper we introduced the elliptic Sombor matrix and the elliptic Sombor energy of a graph  $G$ . We computed the elliptic Sombor characteristic polynomial and the elliptic Sombor energy for some graph classes. Also, we studied the elliptic Sombor energy of cubic graphs of order 10.

## Funding

This research received no external funding.

## Data Availability Statement

Data is contained within the article.

## Conflicts of Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## References

- [1] S. Alikhani, N. Ghanbari, Randić energy of specific graphs, *Appl. Math. Comp.* 269 (2015) 722–730. <https://doi.org/10.1016/j.amc.2015.07.112>
- [2] S. Alikhani, N. Ghanbari, Sombor index of polymers, *MATCH Commun. Math. Comput. Chem.* 86 (2021) 715–728. <https://doi.org/10.48550/arXiv.2103.13663>
- [3] S. B. Bozkurt, D. Bozkurt, Sharp upper bounds for energy and Randić Energy, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 669–680. [https://match.pmf.kg.ac.rs/Match70/n2/match70n2\\_669-680.pdf](https://match.pmf.kg.ac.rs/Match70/n2/match70n2_669-680.pdf)
- [4] S. B. Bozkurt, I. Gutman, Estimating the Incidence Energy, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 143–156. [https://match.pmf.kg.ac.rs/Match70/n1/match70n1\\_143-156.pdf](https://match.pmf.kg.ac.rs/Match70/n1/match70n1_143-156.pdf)
- [5] S. B. Bozkurt, A. D. Güngör, I. Gutman, A. S. Cevik, Randić matrix and Randić energy, *MATCH Commun. Math. Comput. Chem.* 64 (2010) 239–250. <https://www.researchgate.net/publication/286853985>
- [6] H. Chen, W. Li, J. Wang, Extremal values on the Sombor index of trees, *MATCH Commun. Math. Comput. Chem.* 87 (2022) 23–49. <https://doi.org/10.46793/match.87-1.023C>
- [7] L. Chen, Y. Shi, Maximal matching energy of tricyclic graphs, *MATCH Commun. Math. Comput. Chem.* 73 (2015) 105–119. [https://match.pmf.kg.ac.rs/Match73/n1/match73n1\\_105-119.pdf](https://match.pmf.kg.ac.rs/Match73/n1/match73n1_105-119.pdf)
- [8] R. Cruz, I. Gutman and J. Rada, Sombor index of chemical graphs, *Appl. Math. Comp.* 399 (2021) 126018. <https://doi.org/10.1016/j.amc.2021.126018>
- [9] D. Cvetković, M. Doob, H. Sachs, *Spectra of graphs - Theory and Application*, Academic Press, New York, 1980. <https://d-nb.info/942237862/04>
- [10] K. C. Das, A. S. Cevik, I. N. Cangul, Y. Shang, On Sombor index, *Symmetry* 13 (2021) 140. <https://doi.org/10.3390/sym13010140>
- [11] K. C. Das, I. Gutman, A. S. Cevik, B. Zhou, On Laplacian Energy, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 689–696. [https://match.pmf.kg.ac.rs/Match70/n2/match70n2\\_689-696.pdf](https://match.pmf.kg.ac.rs/Match70/n2/match70n2_689-696.pdf)
- [12] K. C. Das, S. Sorgun, On Randić Energy of Graphs, *MATCH Commun. Math. Comput. Chem.* 72 (2014) 227–238. [https://match.pmf.kg.ac.rs/Match72/n1/match72n1\\_227-238.pdf](https://match.pmf.kg.ac.rs/Match72/n1/match72n1_227-238.pdf)
- [13] H. Deng, Z. Tang, R. Wu, Molecular trees with extremal values of Sombor indices, *Int. J. Quantum Chem.* <https://doi.org/10.1002/qua.26622>
- [14] C. Espinal, I. Gutman, J. Rada, Elliptic Sombor index of chemical graphs, *Commun. Comb. Optim.* (2024) 1–11. <https://doi.org/10.22049/cco.2024.29404.1977>
- [15] G. B. Khosrovshahi, Ch. Maysoori, Tayfeh-Rezaie, A Note on 3-Factorizations of  $K_{10}$ , *J. Combin. Designs* 9 (2001) 379–383. <https://doi.org/10.1002/jcd.1018>
- [16] N. Ghanbari, On the Sombor characteristic polynomial and Sombor energy of a graph, *Comp. Appl. Math.* 41 242 (2022). <https://doi.org/10.1007/s40314-022-01957-5>

- [17] N. Ghanbari, S. Alikhani, Sombor index of certain graphs, Iranian J. Math. Chem. 12(1) (2021) 27–37. <https://doi.org/10.22052/ijmc.2021.242106.1547>
- [18] K. J. Gowtham, N. N. Swamy, On Sombor energy of graphs, Nanosystems: Physics, Chemistry, Mathematics 12(4) (2021) 411–417. <https://doi.org/10.17586/2220-8054-2021-12-4-411-417>
- [19] I. Gutman, Geometric approach to degree based topological indices, MATCH Commun. Math. Comput. Chem. 86(1) (2021) 11–16. <https://doi.org/10.1002/qua.27346>
- [20] I. Gutman, Spectrum and energy of the Sombor matrix, Vojnotehnicki Glasnik. 69(3) (2021) 551–561. <https://doi.org/10.5937/vojtehg69-31995>
- [21] I. Gutman, The energy of a graph: Old and new results, in: A. Betten, A. Kohnert, R. Laue, A. Wassermann (Eds.), Algebraic Combinatorics and Applications, Springer-Verlag, Berlin (2001) 196–211. [https://doi.org/10.1007/978-3-642-59448-9\\_13](https://doi.org/10.1007/978-3-642-59448-9_13)
- [22] I. Gutman, Topology and stability of conjugated hydrocarbons. The dependence of total  $\pi$ -electron energy on molecular topology, J. Serb. Chem. Soc. 70 (2005) 441–456. <https://doi.org/10.2298/JSC0503441G>
- [23] I. Gutman, B. Furtula, S. B. Bozkurt, On Randić energy, Linear Algebra Appl. 442 (2014) 50–57. <https://doi.org/10.1016/j.laa.2013.06.010>
- [24] I. Gutman, X. Li, J. Zhang, Graph energy, in: M. Dehmer, F. Emmert-Streib (Eds.), Analysis of Complex Networks. From Biology to Linguistics, Wiley-VCH, Weinheim, (2009) 145–174. <https://doi.org/10.1002/9783527627981.ch7>
- [25] I. Gutman, I. Redžepović, Sombor energy and Hückel rule, Discrete Math. Lett. 9 (2022) 67–71. <https://doi.org/10.47443/dml.2021.s211>
- [26] G. K. Jayanna, I. Gutman, On characteristic polynomial and energy of Sombor matrix, Open J. Discrete Appl. Math. 4(3) (2021) 29–35. <https://doi.org/10.30538/psrp-odam2021.0062>
- [27] S. Ji, X. Li, Y. Shi, Extremal matching energy of bicyclic graphs, MATCH Commun. Math. Comput. Chem. 70 (2013) 697–706. <https://www.researchgate.net/publication/253241318>
- [28] S. Li, Z. Wang, M. Zhang, On the extremal Sombor index of trees with a given diameter, Appl. Math. Comput. 416 (2022) 126731. <https://doi.org/10.1016/j.amc.2021.126731>
- [29] S. Majstorović, A. Klobučar, I. Gutman, Selected topics from the theory of graph energy: hypoenergetic graphs, in: D. Cvetković, I. Gutman (Eds.), Applications of Graph Spectra, Math. Inst., Belgrade, (2009) 65–105. <http://elib.mi.sanu.ac.rs/files/journals/zr/21/n021p065.pdf>
- [30] M. Mohammadi, H. Barzegar, Bounds for Sombor index using topological and statistical indices, J. Disc. Math. Appl. 10(1) (2025) 61–85. <https://doi.org/10.22061/jdma.2025.11494.1107>
- [31] I. Redžepović, Chemical applicability of Sombor indices, J. Serb. Chem. Soc. 86 (2021) 445–457. <https://doi.org/10.2298/JSC201215006R>
- [32] S. Rouhani, M. Habibi, M. A. Mehrpouya, On Sombor index of extremal graphs, J. Disc. Math. Appl. 9(4) (2024) 335–344. <https://doi.org/10.22061/jdma.2024.11328.1101>
- [33] D. Stevanović, M. Milošević, P. Hic, M. Pokorný, Proof of a conjecture on distance energy of complete multipartite graphs, MATCH Commun. Math. Comput. Chem. 70 (2013) 157–162. <https://doi.org/10.1016/j.laa.2020.10.029>
- [34] Z. Wang, Y. Mao, Y. Li, B. Furtula, On relations between Sombor and other degree-based indices, J. Appl. Math. Comput. 68 (2022) 1–17. <https://doi.org/10.1007/s12190-021-01516-x>

**Citation:** S. Alikhani, N. Ghanbari, M. A. Dehghanizadeh, Elliptic Sombor energy of a graph, J. Disc. Math. Appl. 10(2) (2025) 26–38.

 <https://doi.org/10.22061/jdma.2024.11190.1089>



#### COPYRIGHTS

©2024 The author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.