



On the energy of fullerene graphs

Mahin Songhori, Modjtaba Ghorbani *

Department of Mathematics, Faculty of Science, Shahid Rajaei Teacher Training University, Tehran, 16785-136, I. R. Iran

Academic Editor: Ali Reza Ashrafi

Abstract. The concept of energy of graph is defined as the sum of the absolute values of the eigenvalues of a graph. Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be eigenvalues of graph G , the energy of G is defined as $\mathcal{E}(G) = \sum_{i=1}^n |\lambda_i|$. The aim of this paper is to compute the eigenvalues of two fullerene graphs C_{60} and C_{80} .

Keywords. eigenvalue, fullerene, graph energy.

1 Introduction

By using method of graph theory, we can construct cubic graphs whose faces are pentagons and hexagons. We call these graphs fullerene graphs. The fullerene era started in 1985 by Kroto and his co-authors with the discovery of a stable cluster C_{60} and its interpretation as a cage structure with the familiar shape of a soccer ball, see [22]. The well-known fullerene, the C_{60} molecule (see Figure 1), is a closed-cage carbon molecule with three-coordinate carbon atoms tiling the spherical or nearly spherical surface with a truncated icosahedral structure formed by 20 hexagonal and 12 pentagonal rings [23]. Let p , h , n and m be the number of pentagons, hexagons, carbon atoms and bonds between them, in a given fullerene F . Since each atom lies in exactly 3 faces and each edge lies in 2 faces, the number of atoms is $n = (5p + 6h)/3$, the number of edges is $m = (5p + 6h)/2 = 3/2n$ and the number of faces is $f = p + h$. By the Euler's formula, $n - m + f = 2$, one can deduce that $(5p + 6h)/3 = (5p + 6h)/2 + p + h = 2$, and therefore, $p = 12$, $v = 2h + 20$ and $e = 3h + 30$. This implies that such molecules made up entirely of n carbon atoms having 12 pentagonal and

*Corresponding author (Email address: mghorbani@sru.ac.ir).




Figure 1. The IPR Fullerene C_{60} .

$(n/2 - 10)$ hexagonal faces, where $n \neq 22$ is a natural number equal or greater than 20. The goal of this paper is to compute some new results on the eigenvalues of fullerene graphs. We encourage the interested readers to consult paper [18]. For more information on this topic and more details about mathematics of fullerene graphs see Ref.s [1–11] as well as [16].

2 Definitions and preliminaries

Now we recall some algebraic definitions that will be used in this paper. Throughout this paper, our notation is standard and mainly taken from [12–15, 17, 21]. Let G be a simple molecular graph without directed and multiple edges and without loops. The vertex and edge-sets of G are represented by $V(G)$ and $E(G)$, respectively. The adjacency matrix $A(G)$ of graph G with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ is the $n \times n$ symmetric matrix $[a_{ij}]$ such that, $a_{ij} = 1$ if v_i and v_j are adjacent and 0 otherwise. The characteristic polynomial of graph G is defined as

$$\chi(G, \lambda) = \det(A(G) - \lambda I).$$

The roots of this polynomial are eigenvalues of G and form the spectrum of graph as follows:

$$\text{spec}(G) = \{[\lambda_1]^{m_1}, \dots, [\lambda_s]^{m_s}\},$$

where m_i is the multiplicity of eigenvalue λ_i . If G is a graph on n vertices and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of its adjacency matrix, then the energy [20] of G is defined as

$$\mathcal{E}(G) = \sum_{i=1}^n |\lambda_i|.$$




Figure 2. The IPR Fullerene C_{70} .

In theoretical chemistry, the energy is a graph parameter stemming from the Hückel molecular orbital approximation for the total π -electron energy. So, the graph energy has some specific chemical interests and has been extensively studied [19].

Example 2.1. Consider the fullerene C_{70} as depicted in Figure 2. This fullerene is one of the most famous member of the fullerenes, since in this graph all pentagons are isolated. In other words, this fullerene obeys in the Isolated Pentagon Rule (**IPR**). This class of fullerenes is the most stable and many of mathematician work on **IPR** fullerenes.

The eigenvalues of this fullerenes are reported in Table 1. Hence, one can see that the energy of this fullerene is

$$\mathcal{E}(C_{70}) = 109.$$

The aim of this paper is to propose a method for computing the energy of fullerene graphs by means of block matrices. Notice that according a result from the seminal paper of Gutman [20], the energy and graph energy for molecules with bipartite molecular graphs are the same, but fullerenes are not bipartite.

3 Main Results

A bijection σ on V with this property that $e = uv$ is an edge if and only if $\sigma(e) = \sigma(u)\sigma(v)$ is an edge of E , is called an automorphism of graph G . The set of all automorphisms of G under the composition of mappings forms a group denoted by $Aut(G)$.

A circulant matrix is a matrix where each row vector is rotated one element to the right relative to the preceding row vector. In other words, a circulant matrix [23] is specified by

one vector c which appears as the first column of C . The remaining columns of C are each cyclic permutations of the vector c with offset equal to the column index. The last row of C is the vector c , in reverse order and the remaining rows are each cyclic permutations of the last row. In general, an $n \times n$ circulant matrix C takes the following form:

$$C = \begin{pmatrix} c_0 & c_{n-1} & \cdots & c_2 & c_1 \\ c_1 & c_0 & c_{n-1} & \cdots & c_2 \\ \vdots & c_1 & c_0 & \ddots & \vdots \\ c_{n-2} & \ddots & \ddots & \ddots & c_{n-1} \\ c_{n-1} & c_{n-2} & \cdots & c_1 & c_0 \end{pmatrix}.$$

The eigenvectors of a circulant matrix are given by

$$v_j = (1, \omega_j, \omega_j^2, \dots, \omega_j^{n-1})^T, \quad j = 0, 1, \dots, n-1,$$

where, $\omega_k = e^{\frac{2k\pi}{n}i}$ are the n -th roots of unity and $i^2 = -1$. The corresponding eigenvalues are then given by

$$\lambda_j = c_0 + c_{n-1}\omega_j + \cdots + c_1\omega_j^{n-1}, \quad j = 0, \dots, n-1.$$

Let A and B be matrices of dimensions $n \times m$ and $n' \times m'$, respectively. Then their tensor product is an $nm' \times mm'$ matrix with block forms

$$A \otimes B = [a_{ij}B].$$

Theorem 3.1. ([24]) Let $A_{ij}, 1 \leq i, j \leq l$ be square matrices of order n that have the complete set of eigenvectors $\{V_1, \dots, V_n\}$ with $A_{ij}V_k = \alpha_{ij}^k V_k$. Let also $B_k = [\alpha_{ij}^k], 1 \leq k \leq n$ be square matrices of order l , each with a complete set of eigenvectors $\{U_1^k, \dots, U_l^k\}$ satisfying $B_k U_j^k = \beta_j^k U_j^k$ for $1 \leq j \leq l$. Then a complete set of eigenvectors $\{W_1, \dots, W_{nl}\}$ for the square matrix

$$\begin{pmatrix} A_{11} & A_{12} & \cdots & A_{1l} \\ A_{21} & A_{22} & \cdots & A_{2l} \\ \vdots & \vdots & \ddots & \vdots \\ A_{l1} & A_{l2} & \cdots & A_{ll} \end{pmatrix}$$

is given by $W_{(k-1)l+j} = U_j^k \otimes V_k$ for $k = 1, 2, \dots, n$ and $j = 1, 2, \dots, l$. The corresponding eigenvalues are $\lambda_{(k-1)l+j} = \beta_j^k$.

We will apply this theorem to the case where all blocks in the adjacency matrix are circulant matrices. An l -level circulant is one whose adjacency matrix has an $l \times l$ block form A , all A_{ij} being circulant. For example, a 2-level circulant,

$$G = C_n(\{n_i^1\}, \{n_i^2\}, \{m_i^{12}\}),$$

would consist of two vertex sets $S_1 = \{v_1, \dots, v_n\}$ and $S_2 = \{w_1, \dots, w_n\}$ such that




Figure 3. Fullerene graph C_{80} with symmetric group I_h .

m	Eigen	m	Eigen	m	Eigen
3	-1.2	4	1.38	1	3
8	-1.62	6	1	3	2.82
5	-1.93	8	0.62	5	2.47
4	-2.2	4	0.27	3	2.08
4	-2.65	4	-0.71	4	1.91
3	-2.7	6	-1	5	1.46

Table 5. Eigenvalues of icosahedral fullerene C_{80} .

fullerene graphs. Estimating the energy of fullerene graphs were also computed in some earlier papers such as [12].

References

- [1] A. R. Ashrafi, M. Ghorbani and M. Jalali, The vertex PI and Szeged indices of an infinite family of fullerenes, *J. Theor. Comput. Chem.* 7 (2008) 221–231.
- [2] A. R. Ashrafi, M. Jalali, M. Ghorbani and M. V. Diudea, Computing PI and Omega polynomials of an infinite family of fullerenes, *MATCH Commun. Math. Comput. Chem.* 60 (2008) 905–916.
- [3] N. Biggs, *Algebraic Graph Theory*, Cambridge Univ. Press, Cambridge, 1974.
- [4] D. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs—Theory and Applications*, Barth, Heidelberg, 1995.
- [5] D. Cvetković, P. Rowlinson, P. Fowler and D. Stevanović, Constructing fullerene graphs from their eigenvalues and angles, *Linear Algebra Appl.* 356 (2002) 37–56.
- [6] E. Estrad, Characterization of 3D molecular structure, *Chem. Phys. Lett.* 319 (2000) 713–718.
- [7] G. H. Fath-Tabar, A. R. Ashrafi and D. Stevanović, Spectral properties of fullerenes, *J. Comput. Theor. Nanosci.* 9 (1) (2012) 327–329.
- [8] P. W. Fowler and D. E. Manolopoulos, *An Atlas of Fullerenes*, Clarendon Press, Oxford, 1995.
- [9] M. Ghorbani and A. R. Ashrafi, Counting the number of hetero fullerenes, *J. Comput. Theor. Nanosci.* 3 (2006) 803–810.
- [10] M. Ghorbani, A. R. Ashrafi and M. Hemmasi, Eccentric connectivity polynomial of C_{18n+10} Fullerenes, *Bulg. Chem. Commun.* 45 (2013) 5–8.
- [11] M. Ghorbani, M. Faghani, A. R. Ashrafi, S. Heidari-Rad and A. Graovać, An upper bound for energy of matrices associated to an infinite class of fullerenes, *MATCH Commun. Math. Comput. Chem.* 71 (2014) 341–354.
- [12] M. Ghorbani and S. Heidari-Rad, Study of fullerenes by their algebraic properties, *Iranian J. Math. Chem.* 3 (2012) 9–24.
- [13] M. Ghorbani and E. Naserpour, Study of some nanostructures by using their Kekulé structures, *J. Comput. Theor. Nanosci.* 10 (2013) 2260–2263.
- [14] M. Ghorbani and M. Songhori, The enumeration of hetero-fullerenes by Polya’s theorem, Fullerenes, Nanotubes and Carbon Nanostructures, *J. Comput. Theor. Nanosci.* 21 (2013) 460–471.
- [15] M. Ghorbani and E. Bani-Asadi, Remarks on characteristic coefficients of fullerene graphs, *Appl. Math. Comput.* 230 (2014) 428–435.
- [16] M. Ghorbani, Remarks on markaracter table of fullerene graphs, *J. Comput. Theor. Nanosci.* 11 (2014) 363–379.
- [17] C. Godsil and G. Royle, *Algebraic Graph Theory*, Springer-Verlag, New York, 2001.
- [18] A. Graovać, O. Ori, M. Faghani and A. R. Ashrafi, Distance property of fullerenes, *Iranian J. Math. Chem.* 2 (2011) 99–107.
- [19] I. Gutman and O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, Berlin, 1986.
- [20] I. Gutman, The energy of a graph, *Ber. Math.-Statist. Sect. Forsch. Graz* 103 (1978) 1–22.
- [21] F. Harary, *Graph Theory*, Addison-Wesley, Reading, MA, 1969.
- [22] H. W. Kroto, J. R. Heath, S. C. ÓBrien, R. F. Curl and R. E. Smalley, buckminster fullerene, *Nature* 318 (1985) 162–163.
- [23] H. W. Kroto, J. E. Fichier and D. E. Cox, *The Fullerene*, Pergamon Press, New York 1993.
- [24] S. L. Lee, Y. L. Luo, B. E. Sagan and Y. -N. Yeh, Eigenvectors and eigenvalues of some special graphs, IV multilevel circulants, *Int. J. Quant. Chem.* 41 (1992) 105–116.
- [25] W. C. Shiu, On the spectra of the fullerenes that contain a nontrivial cyclic-5-cutset, *Australian J. Combin.* 47 (2010) 41–51.