Journal of Discrete Mathematics and Its Applications 7 (2) (2022) 73–80

Journal of Discrete Mathematics and Its Applications

Available Online at: http://jdma.sru.ac.ir

Shahid Rajaee Teacher Training University

On borderenergetic and L-borderenergetic graphs

Mardjan Hakimi-Nezhaad *

Department of Mathematics, Faculty of Science, Shahid Rajaee Teacher Training University, Tehran, 16785 - 163, I. R. Iran

Academic Editor: Mahin Songhori

Abstract. A graph G of order n is said to be borderenergetic if its energy is equal to $2n-2$. In this paper, we study the borderenergetic and Laplacian borderenergetic graphs.

Keywords: energy (of graph), adjacency matrix, Laplacian matrix, signless Laplacian matrix
Mathematics Subject Classification (2010): 05C76.

1 Introduction

We first recall some definitions that will be kept throughout. Let G be a simple graph with n vertices and $m(G)$ edges, and $A(G)$ denotes its adjacency matrix. Let $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$ be the Laplacian and signless Laplacian matrix of the graph G, respectively, where $D(G) = [d_{ij}]$ is the diagonal matrix whose entries are the degree of vertices, i.e., $d_{ii} = deg(v_i)$ and $d_{ij} = 0$ for $i \neq j$.

The energy of G is a graph invariant which was introduced by Ivan Gutman [6]. It is defined as $E(G) = \sum_{i=1}^{n} |\lambda_i|$, where λ_i's are eigenvalues of G. If $0 = \mu_1 \leq \mu_2 \leq \cdots \leq \mu_{n-1} \leq \mu_n$ and $q_1 \leq q_2 \leq \cdots \leq q_{n-1} \leq q_n$ are the Laplacian and signless Laplacian eigenvalues of G then the quantities $E_L(G) = \sum_{i=1}^{n} |\mu_i - \frac{2m(G)}{n}|$ and $E_Q(G) = \sum_{i=1}^{n} |q_i - \frac{2m(G)}{n}|$ are called the Laplacian and signless Laplacian energy of G, respectively. Details on the properties of Laplacian and signless Laplacian energy can be found in [6, 8, 13].

Email address: m.hakiminezhaad@sru.ac.ir
Received 14 April 2022; Revised 28 April 2022; Accepted 14 May 2022
First Publish Date: 1 June 2022

 ©Shahid Rajaee Teacher Training University

The first borderenergetic graph was discovered by Hou et al. in 2001 [11], but in that time it did not attract much attention. Recently, Gong et al. in [5] studied the graphs with the same energy as a complete graph. They put forward the concept of borderenergetic graphs.

A graph G on n vertices is said to be borderenergetic if its energy equals the energy of the complete graph K_n, i.e., if $E(G) = E(K_n) = 2(n-1)$. In [5] it was shown that there exist borderenergetic graphs on order n for each integer $n \geq 7$. The number of borderenergetic graphs were determined for $n = 7,8,9$ [5], $n = 10,11$ [14,17] and $n = 12$ [4].

In [12] a family of non-regular and non-integral borderenergetic threshold graphs was discovered. In [3] the authors obtained three asymptotically tight bounds on the number of edges of borderenergetic graphs. We refer the readers to [9,15] for more information.

An analogous concept as borderenergetic graphs, called Laplacian borderenergetic graphs was proposed in [19]. That is, a graph G of order n is Laplacian borderenergetic or L-border energetic for short, if $E_L(G) = E_L(K_n) = 2n-2$.

In [1] Deng et al. presented some asymptotically bounds on the order and size of L-borderenergetic graphs. Also, they showed that all trees, cycles, the complete bipartite graphs, and many 2-connected graphs are not L-borderenergetic. They showed in [2] a kind of threshold graphs are L-borderenergetic.

Lu et al. in [16] presented all non-complete L-borderenergetic graphs of order $4 \leq n \leq 7$ and they constructed one connected non-complete L-borderenergetic graph on n vertices for each integer $n \geq 4$, which extends the result in [20] and completely confirms the existence of non-complete L-borderenergetic graphs. Particularly, they proved that there are at least $\frac{n}{2}+4$ non-complete L-borderenergetic graphs of order n for any even integer $n \geq 6$.

Hakimi-Nezhaad et al. in [10] generalized the concept of borderenegetic graphs for the signless Laplacian matrices of graphs. That is, a graph G of order n is signless Laplacian borderenergetic or Q-borderenergetic for short, if $E_Q(G) = E_Q(K_n) = 2n-2$. Also, they constructed sequences of Laplacian borderenergetic non-complete graphs by means of graph operations, and all the non-complete and pairwise non-isomorphic L-borderenergetic and Q-borderenergetic graphs of small order n are depicted for n with $4 \leq n \leq 9$, see Appendix.

Tao et al. in [18] considered the extremal number of edges of non-complete L-borderenergetic graph, then use a computer search to find out all the L-borderenergetic graphs on no more than 10 vertices.

Main results

Here, we present some basic theorem used to study borderenergetic and L-borderenergetic and Q-borderenergetic graphs.

Theorem 1.1. *We have the following statements:*

1) [5]. *There are no borderenergetic graphs of order* $n \leq 6$.

2) [5]. *There exists a unique borderenergetic graph of order 7.*

3) [5]. *For any* $n \geq 7$, *there exist borderenergetic graphs of order* n.

4) [5]. *There are exactly 6 borderenergetic graphs of order 8.*

5) [5]. *There are exactly 17 borderenergetic graphs of order 9.*

6) [14,17]. *There are exactly 49 borderenergetic graphs of order 10.*

7) [17]. *There are exactly 158 borderenergetic graphs of order 11, of which 157 are connected.*

8) [4]. *There are exactly 572 connected borderenergetic graphs of order 12.*

8) [5]. *For each integer* n $(n \geq 13)$, *there exists a non-complete borderenergetic graph of order* n.

Theorem 1.2. [9]. *A borderenergetic graph of order* n *must possess at least* $2n - 2$ *edges.*

Theorem 1.3. [3]. *Let* G *be a* k*-regular integral graph of order* n *with* t *non-negative eigenvalues. If* $E(G) = 2(n - t + k)$ *then* $E(\bar{G}) = 2(n - 1)$, *where* $\bar{G}$ *is complement of graph* G.

Theorem 1.4. [15].

1) *There is no noncomplete borderenergetic graph with maximum degree* $\triangle = 2$ *or* 3.

2) *Let* G *be a non-complete borderenergetic graph of order* n *with maximum degree* $\triangle = 4$. *Then* G *must have the following properties:*

(i) $e(G) = 2n$ *or* $2n - 1$;

(ii) $|G| \leq 21$;

(iii) G *is non-bipartite;*

(iv) the nullity, i.e., the multiplicity of eigenvalue 0, of G *is 0.*

3) *Let* G *be a* 4*-regular non-complete borderenergetic graph of order* n *and* H *is a maximal bipartite subgraph of* G. *Then* $m(G) - m(H) \geq 3$.

Theorem 1.5. [15]. *No borderenergetic graphs have minimum degree* $n - 2$. *Besides, for each integer* $n \geq 7$, *there exists a connected noncomplete borderenergetic graph of order* n *with minimum degree* $n - 3$ *and for each even integer* $n \geq 8$, *there exists a noncomplete borderenergetic graph of order* n *with minimum degree* $n - 4$.

Theorem 1.6. *We have the following statements:*

1) [10]. *There are exactly two non-complete L-borderenergetic disconnected graphs of orders 4 and 5, respectively.*

2) [10]. *There are exactly five non-complete L-borderenergetic disconnected graphs of order 6.*

3) [10]. *There are exactly five non-complete L-borderenergetic disconnected graphs of order 7.*

4) [18]. *There are totally 18 L-borderenergetic connected graphs on less than 8 vertices.*

5) [10,18]. *There are exactly 31 L-borderenergetic connected graphs and 27 disconnected graphs of order 8.*

6) [10,18]. *There are exactly 16 L-borderenergetic graphs and 26 disconnected graphs of order 9.*

7) [10,18]. *There are exactly 120 L-borderenergetic connected graphs on 10 vertices.*

Theorem 1.7. [10] *There is no Laplacian borderenergetic tree with $n \geq 3$ vertices.*

Theorem 1.8. [1]. *If G is a complete bipartite graph $K_{a,b}(1F\text{œ}F\text{ø})$, then G is not L-borderenergetic.*

Theorem 1.9. [1]. *If G is a 2-connected graph with maximum degree $\triangle = 3$ and $t(G) \geq 7$ then G is not L-borderenergetic, where $t(G)$ the number of vertices of degree 3 in G.*

Theorem 1.10. [10].

1) *There are no non-complete Q-borderenergetic graph of order $n \leq 5$ and 7.*

2) *There are exactly two non-complete Q-borderenergetic of order 6.*

3) *There exist exactly fourteen non-complete Q-borderenergetic graphs of order 8.*

4) *There exist exactly sixteen non-complete Q-borderenergetic graphs of order 9.*

References

[1] B. Deng, X. Li, J. Wang, Further results on L-borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 77 (3) (2017) 607–616.
[2] B. Deng, X. Li, More on L-borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 77 (1) (2017) 115–127.
[3] B. Deng, X. Li, I. Gutman, More on borderenergetic graphs, Linear Algebra Appl. 497 (2016) 199–208.
[4] B. Furtula, I. Gutman, Borderenergetic Graphs of Order 12, Iranian J. Math. Chem. 8 (4) (2017) 339–343.
[5] S. Gong, X. Li, G. Xu, I. Gutman, B. Furtula, Borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 74 (2015) 321–332.
[6] I. Gutman, The energy of a graph, Ber. Math-Statist. Sekt. Forschungszentrum Graz 103 (1978) 1–22.
[7] I. Gutman, The energy of a graph: old and new results, in: A. Betten, A. Kohner, R. Laue, A. Wassermann (Eds.), Algebraic Combinatorics and Applications, Springer, Berlin (2001) 196–211.
[8] I. Gutman, B. Zhou, Laplacian energy of a graph, Linear Algebra Appl. 414 (2006) 29–37.
[9] I. Gutman, On Borderenergetic graphs, Bull. Acad. Serbe Sci. Arts. (Cl. Math. Natur.) 42 (2017) 9–18.
[10] M. Hakimi-Nezhaad, M. Ghorbani, Laplacian Borderenergetic graphs, Journal of Information and Optimization Sciences, inpress.

[11] Y. Hou, I. Gutman, Hyperenergetic line graphs, MATCH Commun. Math. Comput. Chem. 43 (2001) 29–39.
[12] Y. Hou, Q. Tao, Borderenergetic threshold graphs, MATCH Commun. Math. Comput. Chem. 75 (2016) 253–262.
[13] X. Li, Y. Shi, I. Gutman, Graph energy, Springer, New York, 2012.
[14] X. Li, M. Wei, S. Gong, A computer search for the borderenergetic graphs of order 10, MATCH Commun. Math. Comput. Chem. 74 (2015) 333–342.
[15] X. Li, M. Wei, X. Zhu, Borderenergetic graphs with small maximum or large minimum degrees, MATCH Commun. Math. Comput. Chem. 77 (2016) 25–36.
[16] L. Lu, Q. Huang, On the existence of non-complete $L-$borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 77(3) (2017) 625–634.
[17] Z. Shao, F. Deng, Correcting the number of borderenergetic graphs of order 10, MATCH Commun. Math. Comput. Chem. 75 (2016) 263–265.
[18] Q. Tao, Y. Hou, A computer search for the L-borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 77(3) (2017) 595–606.
[19] F. Tura, L-borderenergetic graphs, MATCH Commun. Math. Comput. Chem. 77 (2017) 37–44.
[20] F. Tura, L-borderenergetic graphs and normalized Laplacian energy, MATCH Commun. Math. Comput. Chem. 77(3) (2017) 617–624.

Appendix




Figure 1. All Laplacian borderenergetic dis-connected graphs of order 4 and 5.




Figure 2. All Laplacian borderenergetic dis-connected graphs of order 6.




Figure 3. All Laplacian borderenergetic dis-connected graphs of order 7.




Figure 4. All Laplacian borderenergetic dis-connected graphs of order 8.




Figure 5. All Laplacian borderenergetic dis-connected graphs of order 9.

Citation: M. Hakimi-Nezhaad, On borderenergetic and L-borderenergetic graphs, J. Disc. Math. Appl. 7(2) (2022) 73–80.
https://doi.org/10.22061/jdma.2022.513



COPYRIGHTS
©2023 The author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.